

ABSTRACT ELEMENTARY CLASSES AND ACCESSIBLE CATEGORIES

T. BEKE AND J. ROSICKÝ*

ABSTRACT. We compare abstract elementary classes of Shelah with accessible categories having directed colimits.

1. INTRODUCTION

Infinitary model theory deals with the properties of models of infinitary first-order theories. M. Makkai and R. Paré [10] initiated a category-theoretic investigation of the subject by introducing *accessible categories* and *accessible functors*. Accessible categories are categories of models $\mathbf{Mod}(T)$ of suitable theories T in a language $L_{\kappa\lambda}(\Sigma)$ where Σ is a signature. Morphisms in $\mathbf{Mod}(T)$ are homomorphisms. Makkai and Paré found purely categorical characterizations of categories of this form and showed, among many other facts, that for every theory T of $L_{\kappa\lambda}(\Sigma)$, $\mathbf{Elem}(T)$, the category of T -models together with $L_{\kappa\lambda}$ -elementary embeddings forms an accessible category. Abstract elementary classes were introduced by S. Shelah with a similar aim – to formalize properties of models of infinitary first-order theories of finitary signatures (see [2]), especially the cardinalities of non-isomorphic models, without explicit reference to syntax. There is a visible connection of abstract elementary classes with accessible categories; we realized this in 2007. Around the same time, M. Lieberman arrived at very similar results, characterizing abstract elementary classes as suitable subcategories of categories $\mathbf{Emb}(\Sigma)$ of Σ -structures and submodel embeddings where Σ is finitary; see [8] and [9]. The role of this ambient category can also be played by $\mathbf{Str}(\Sigma)$, the category of Σ -structures and homomorphisms, and by other finitely accessible categories.

Unlike abstract elementary classes, accessible categories are not equipped with canonical ‘underlying sets’. Nonetheless, a good substitute for ‘size of the model’, namely, the *presentability rank* of an object, can be expressed purely in the language of category theory, i.e. in terms of

Date: May 11, 2010.

Key words and phrases. abstract elementary class, accessible category.

* Supported by MSM 0021622409.

objects (models) and morphisms (embeddings). In this sense, accessible categories take an extreme ‘signature-free’ and ‘elements-free’ view of abstract elementary classes (AEC). From this point of view, Shelah’s Categoricity Conjecture, the driving force of AEC (see [2]), turns on the subtle interaction between ranks of objects and directed colimits in accessible categories.

Abstract elementary classes turn out to be placed between $\mathbf{Elem}(T)$ where T is a theory of $L_{\kappa\omega}$ and general accessible categories with directed colimits whose morphisms are monomorphisms.

2. ACCESSIBLE CATEGORIES

In order to define accessible categories one just needs the concept of a λ -directed colimit where λ is a regular cardinal number. This is a colimit over a diagram $D : \mathcal{D} \rightarrow \mathcal{K}$ where \mathcal{D} is λ -directed poset (taken as a category). An object K of a category \mathcal{K} is called λ -presentable if its hom-functor $\text{hom}(K, -) : \mathcal{K} \rightarrow \mathbf{Set}$ preserves λ -directed colimits; \mathbf{Set} is the category of sets.

A category \mathcal{K} is called λ -accessible, where λ is a regular cardinal, provided that

- (1) \mathcal{K} has λ -directed colimits,
- (2) \mathcal{K} has a set \mathcal{A} of λ -presentable objects such that every object of \mathcal{K} is a λ -directed colimit of objects from \mathcal{A} .

A category is *accessible* if it is λ -accessible for some regular cardinal λ .

Definition 2.1. We say that a category \mathcal{K} is *well λ -accessible* if it is μ -accessible for each regular cardinal $\lambda \leq \mu$.

\mathcal{K} is *well accessible* if it is well λ -accessible for some regular cardinal λ .

Remark 2.2. (i) *Locally presentable categories* are defined as cocomplete accessible categories. Following [1], 1.20, each locally λ -presentable category is well λ -accessible.

(ii) Let λ be an uncountable regular cardinal. The category \mathbf{Pos}_λ of λ -directed posets and order embeddings is λ -accessible but it is not well accessible. In fact, \mathbf{Pos}_λ is μ -accessible if and only if $\lambda \triangleleft \mu$. More generally, if $\lambda \triangleleft \mu$ then any λ -accessible category \mathcal{K} is μ -accessible, but a λ -accessible category may fail to be μ -accessible for a proper class of regular cardinals $\lambda < \mu$. See [10], 2.3.10 or [1], 2.11.

(iii) Since $\omega \triangleleft \lambda$ for each uncountable regular cardinal λ (see [1], 2.13 (1)), each finitely accessible category is well finitely accessible.

A *signature* Σ is a set of (infinitary) relation and function symbols. These symbols are S -sorted where S is a set of sorts. It is advantageous

to work with many-sorted signatures but it is easy to reduce them to single-sorted ones. One just replaces sorts by unary relation symbols and adds axioms saying that there are disjoint and cover the underlying set of a model. Thus the underlying set of an S -sorted structure A is the disjoint union of underlying sets A_s over all sorts $s \in S$. The category of all Σ -structures and homomorphisms is denoted by $\mathbf{Str}(\Sigma)$ and the category of all Σ -structures and submodel embeddings by $\mathbf{Emb}(\Sigma)$. Both $\mathbf{Str}(\Sigma)$ and $\mathbf{Emb}(\Sigma)$ are accessible categories, cf. [1], 5.30 and 1.70.

A signature Σ is *finitary* if all relation and function symbols are finitary. For a finitary signature, the category $\mathbf{Str}(\Sigma)$ is finitely accessible and $\mathbf{Emb}(\Sigma)$ is well accessible and has directed colimits. both cases, there is a cardinal κ such that, for each regular cardinal $\kappa \leq \mu$, a Σ -structure K is μ -presentable if and only if $|K| < \mu$. This follows from the downward Löwenheim-Skolem theorem ([10], 3.3.1).

Example 2.3. Let Σ be one-sorted with countably many unary relation symbols R_n , $n = 0, 1, \dots, n, \dots$. Let $A = \{a\}$ be the Σ -structure on a singleton set $\{a\}$ that satisfies all relations R_n . Then A is not finitely presentable, and does not have any non-trivial finitely presentable submodels. Thus $\mathbf{Emb}(\Sigma)$ is not finitely accessible. It is, however, ω_1 -accessible.

Let λ be a cardinal and Σ be a λ -ary signature, i.e., all relation and function symbols have arity smaller than λ . Given a cardinal κ , the language $L_{\kappa\lambda}(\Sigma)$ allows less than κ -ary conjunctions and disjunctions and less than λ -ary quantifications. A *theory* T is a set of sentences of $L_{\kappa\lambda}(\Sigma)$. $\mathbf{Mod}(T)$ denotes the category of T -models and homomorphisms, $\mathbf{Emb}(T)$ the category of T -models and submodel embeddings while $\mathbf{Elem}(T)$ will denote the category of T -models and $L_{\kappa\lambda}$ -elementary embeddings. The category $\mathbf{Elem}(T)$ is accessible (see [1], 5.42). For certain theories T , the category $\mathbf{Mod}(T)$ does not have μ -directed colimits for any regular cardinal μ and thus fails to be accessible.

A theory T is called *basic* if it consists of sentences

$$\forall(x)(\varphi(x) \Rightarrow \psi(x))$$

where φ and ψ are positive-existential formulas and x is a string of variables. For a basic theory T , the category $\mathbf{Mod}(T)$ is accessible. Conversely, every accessible category is equivalent to the category of models and homomorphisms of a basic theory. All these facts can be found in [10] or [1]. Let us add that, prior to [10], they were announced in [11].

A functor $F : \mathcal{K} \rightarrow \mathcal{L}$ is called λ -accessible if \mathcal{K} and \mathcal{L} are λ -accessible categories and F preserves λ -directed colimits. F is called accessible if it is λ -accessible for some regular cardinal λ . By the Uniformization theorem (see [10], 2.5.1 or [1], 2.19), for each accessible functor F there is a regular cardinal λ such that F is λ -accessible and preserves λ -presentable objects; it means that if K is λ -presentable then $F(K)$ is λ -presentable. The same is then true for each $\mu \triangleright \lambda$.

Definition 2.4. A functor $F : \mathcal{K} \rightarrow \mathcal{L}$ will be called *well λ -accessible* if \mathcal{K} and \mathcal{L} are well λ -accessible categories and F preserves λ -directed colimits and μ -presentable objects for each $\lambda \leq \mu$. F is called *well accessible* if it is well λ -accessible for some regular cardinal λ .

Remark 2.5. (i) Every colimit preserving functor between cocomplete λ -accessible categories is well λ -accessible. It immediately follows from the fact that, for regular cardinals $\lambda \leq \mu$, an object of a locally λ -presentable category is μ -presentable if and only if it is a μ -small colimit of λ -presentable objects. Recall that a category is μ -small if it has less than μ morphisms.

(ii) Every finitely accessible functor is well finitely accessible. It follows from [1], 2.13 (1) and the fact that, for regular cardinals $\lambda \triangleleft \mu$, an object of a λ -presentable category is μ -presentable if and only if it is a λ -directed μ -small colimit of λ -presentable objects (see [10], 2.3.11).

We say that a functor $F : \mathcal{K} \rightarrow \mathcal{L}$ *reflects λ -presentable objects* if $F(K)$ λ -presentable implies that K is λ -presentable.

Recall that a morphism $g : B \rightarrow A$ is a split epimorphism if there exists $f : A \rightarrow B$ with $gf = \text{id}_A$. Since, in this case, g is a coequalizer of fg and id_B , A is λ -presentable whenever B is λ -presentable (see [1], 1.16).

Definition 2.6. We say that a functor $F : \mathcal{K} \rightarrow \mathcal{L}$ *reflects split epimorphisms* if f is a split epimorphism whenever $F(f)$ is a split epimorphism.

Lemma 2.7. *Let $F : \mathcal{K} \rightarrow \mathcal{L}$ be a λ -accessible functor which reflects split epimorphisms. Then F reflects λ -presentable objects.*

Proof. Let $F(K)$ be λ -presentable in \mathcal{L} . Since \mathcal{K} is λ -accessible, K is a λ -directed colimit $(k_i : K_i \rightarrow K)_{i \in I}$ of λ -presentable objects. Since F preserves λ -directed colimits and $F(K)$ is λ -presentable, there is $i \in I$ and $f : F(K) \rightarrow F(K_i)$ such that $F(k_i)f = \text{id}_{F(K)}$. Since F reflects split epimorphisms, there is $k_i : K_i \rightarrow K$ is a split epimorphism. Thus K is λ -presentable. \square

Definition 2.8. A subcategory \mathcal{K} of a category \mathcal{L} will be called *nearly full* if for each commutative triangle

$$\begin{array}{ccc} A & \xrightarrow{h} & C \\ & \searrow f & \nearrow g \\ & B & \end{array}$$

h and g in \mathcal{K} imply that f is in \mathcal{K} .

Remark 2.9. [6] calls such subcategories coherent. The embedding of a nearly full subcategory \mathcal{K} in \mathcal{L} reflects split epimorphisms.

Definition 2.10. Let λ be a regular cardinal. We say that an object K of a category \mathcal{K} has *presentability rank* λ if it is λ -presentable but not μ -presentable for any regular cardinal $\mu < \lambda$. We will write $\text{rank}(K) = \lambda$.

This concept was introduced by Makkai and Paré (see the very last line of p.29 of [10]) under the name of *presentability* of an object.

Remark 2.11. (1) We have mentioned that in $\mathbf{Emb}(\Sigma)$, Σ finitary, there is a cardinal κ such that, for each cardinal $\kappa \leq \mu$, a Σ -structure A is μ -presentable if and only if $|A| < \mu$. Therefore for all large enough structures, $\text{rank}(A) = |A|^+$, i.e., presentability ranks play the role of cardinalities.

(2) Assuming GCH, in any accessible category \mathcal{K} , $\text{rank}(K)$ is a successor cardinal for all objects K (with the possible exception of a set of isomorphism types). Indeed, let \mathcal{K} be a κ -accessible category and $\text{rank}(K) = \lambda$ with $\kappa < \lambda$. λ is a regular cardinal by definition; if λ was a limit cardinal then (as it is uncountable) it would be a weakly inaccessible cardinal. Since GCH is assumed, λ is inaccessible. Thus, given $\alpha < \kappa < \lambda$ and $\beta < \lambda$, we have $\beta^\alpha < \lambda$. Following [1] 2.13 (4), $\kappa \triangleleft \lambda$. Following [10], 2.3.11, the object K can be exhibited as a κ -directed colimit of κ -presentable objects along a diagram of size less than λ . Let the size of that diagram be ν ; then K is ν^+ -presentable. That would mean $\nu^+ = \lambda$, contradicting that λ is a limit cardinal.

Proposition 2.12. Let $F : \mathcal{K} \rightarrow \mathcal{L}$ be a well λ -accessible functor which reflects split epimorphisms. Then F preserves presentability ranks μ for $\lambda < \mu$.

Proof. Let $\text{rank}(K) = \mu$, $\lambda < \mu$. Then $F(K)$ is μ -presentable. Assume that $F(K)$ is ν -presentable for some $\nu < \mu$. Without loss of generality, we can assume that $\lambda \leq \nu$. Following 2.7, K is ν -presentable, which is a contradiction. Thus $\text{rank}(F(K)) = \mu$. \square

3. ACCESSIBLE CATEGORIES WITH DIRECTED COLIMITS

Proposition 3.1. *Each accessible category with directed colimits is well accessible.*

Proof. Let \mathcal{K} be a λ -accessible category with directed colimits and consider a regular cardinal $\lambda < \mu$. Given an object K of \mathcal{K} , there is a λ -directed colimit $(a_i : A_i \rightarrow K)_{i \in I}$ of λ -presentable objects A_i . Let \hat{I} be the poset of all directed subsets of I of cardinalities less than μ (ordered by inclusion). Since $\omega \triangleleft \mu$, \hat{I} is μ -directed (see [1], 2.13 and 2.11). For each $M \in \hat{I}$, let B_M be a colimit of a subdiagram indexed by M . Then B_M is μ -presentable and K is a μ -directed colimit of B_M , $M \in \hat{I}$ (see the proof of the last implication in [1], 2.11). \square

Proposition 3.2. *Let \mathcal{K} and \mathcal{L} be accessible categories with directed colimits and $F : \mathcal{K} \rightarrow \mathcal{L}$ a functor preserving directed colimits. Then F is well accessible.*

Proof. There is a regular cardinal λ such that both \mathcal{K} and \mathcal{L} are λ -accessible and F preserves λ -presentable objects (see [1] 2.19). Consider a regular cardinal $\lambda < \mu$ and let K be a μ -presentable object of \mathcal{K} . Following the proof of 3.1, K is a μ -directed colimit of objects B_M where each B_M is a directed colimit of less than μ λ -presentable objects. Since K is μ -presentable, it is a retract of some B_M . Since $F(B_M)$ is a directed colimit of less than μ λ -presentable objects, $F(B_M)$ is μ -presentable in \mathcal{L} . Thus $F(K)$ is μ -presentable in \mathcal{L} as a retract of $F(B_M)$. We have proved that F is well λ -accessible. \square

Recall that a full subcategory \mathcal{K} of a category \mathcal{L} is called accessibly embedded if there is a regular cardinal λ such that \mathcal{K} is closed under λ -directed colimits in \mathcal{L} .

Theorem 3.3. *Accessible categories with directed colimits are precisely reflective and accessibly embedded subcategories of finitely accessible categories.*

Proof. Every reflective and accessibly embedded subcategory of a finitely accessible category \mathcal{L} is accessible (see [1], 2.53) and has directed colimits calculated as reflections of directed colimits in \mathcal{L} . Conversely, let \mathcal{K} be a λ -accessible category with directed colimits. Then \mathcal{K} is equivalent to a full subcategory $\text{Ind}_\lambda(\mathcal{C})$ of $\mathbf{Set}^{\mathcal{C}^{\text{op}}}$ consisting of all λ -directed colimits of hom-functors (see [1], 2.26); \mathcal{C} is a small category. Let $\text{Ind}(\mathcal{C})$ be the full subcategory of $\mathbf{Set}^{\mathcal{C}^{\text{op}}}$ consisting of all directed colimits of hom-functors. Then $\text{Ind}_\lambda(\mathcal{C})$ is closed in $\text{Ind}(\mathcal{C})$ under λ -directed colimits. Given an object X in $\text{Ind}(\mathcal{C})$, we express it as a

directed colimit of hom-functors and take their colimit $F(X)$ in $\text{Ind}_\lambda(\mathcal{C})$ (recall that \mathcal{K} has directed colimits). Clearly, $F(X)$ is a reflection of X in $\text{Ind}_\lambda(\mathcal{C})$. Thus $\text{Ind}_\lambda(\mathcal{C})$ is a reflective subcategory of $\text{Ind}(\mathcal{C})$. \square

Definition 3.4. An accessible category \mathcal{K} will be called λ -LS-accessible if \mathcal{K} has an object of presentability rank μ^+ for each cardinal $\mu \geq \lambda$.

\mathcal{K} is LS-accessible if it is λ -LS-accessible for some cardinal λ .

Remark 3.5. (1) Following 2.11 (1), $\mathbf{Emb}(\Sigma)$ is LS-accessible for each finitary signature Σ and $\text{rank}(X) = |X|^+$ for each Σ -structure X .

(2) Let T be a basic theory in a language $L_{\kappa\omega}(\Sigma)$. Since $\mathbf{Emb}(T)$ is closed under directed colimits in $\mathbf{Emb}(\Sigma)$, $\text{rank}(X) \leq |X|^+$ for each T -model X . Following the downward Löwenheim-Skolem theorem ([10] 3.3.1) there is a cardinal μ such that each T -model is a μ^+ -directed colimit of submodels which are T -models and have cardinality μ . Thus, for $\mu \leq |X|$, we have $\text{rank}(X) = |X|^+$. Moreover, $\mathbf{Emb}(T)$ is LS-accessible.

(3) Given an arbitrary theory T a language $L_{\kappa\omega}(\Sigma)$, we can use Morley's device, to find a conservative extension T^+ of the theory T such that T^+ -homomorphisms are those T -homomorphisms that preserve and reflect the prescribed formulas (see [10] 3.2.8 for details). Consequently, $\mathbf{Elem}(T)$ is LS-accessible and $\text{rank}(X) = |X|^+$ for each T -model X .

Theorem 3.6. *Each locally presentable category is LS-accessible.*

Proof. Let \mathcal{K} is a locally presentable category. Then \mathcal{K} is locally λ -presentable for some regular cardinal λ and, following [1] 5.30, \mathcal{K} is equivalent to $\mathbf{Mod}(T)$ for a limit theory of $L_{\lambda\lambda}(\Sigma)$ where Σ is an S -sorted signature. Thus it suffices to prove that $\mathbf{Mod}(T)$ is LS-accessible.

Let \mathbf{Set}^S denote the category of S -sorted sets. The category \mathbf{Set}^S is locally finitely presentable and, given an S -sorted set $X = (X_s)_{s \in S}$, $\text{rank}(X) = |X|^+$ where $|X|$ is defined as the cardinality of the disjoint union of X_s , $s \in S$. Let $U : \mathcal{K} \rightarrow \mathbf{Set}^S$ denote the forgetful functor. The functor U preserves limits and λ -directed colimits (see the proof of [1] 5.9). Thus U has a left adjoint F ([1] 1.66) which preserves μ -presentable objects for each $\mu \geq \lambda$ ([1] Ex. 1.s(1)). Thus $\text{rank}(FX) \leq \text{rank}(X)$ for each X with $\lambda \leq \text{rank}(X)$. Assume that $\mu = \text{rank}(FX) < \text{rank}(X)$ for X with $\lambda \leq \text{rank}(X)$. Since \mathbf{Set}^S is locally μ -presentable, X is a μ -directed colimit of μ -presentable objects X_i , $i \in I$. Let

$$(u_i : X_i \rightarrow X)_{i \in I}$$

denote a colimit cocone. Since F preserves colimits,

$$(Fu_i : FX_i \rightarrow FX)_{i \in I}$$

is a colimit cocone. Since FX is μ -presentable, there is $j \in I$ and $r : FX \rightarrow FX_j$ such that $F(u_j)r = \text{id}_{FX}$. Hence Fu_j is a split epimorphism. We can assume, without a loss of generality, that $X_s \neq \emptyset$ implies $(X_i)_s \neq \emptyset$ for each $s \in S$. Then each u_i is a split monomorphism and thus each Fu_i is a split monomorphism. Consequently, Fu_j is an isomorphism and $r = (Fu_j)^{-1}$.

Let $\eta : \text{Id} \rightarrow UF$ be the adjunction unit. Then

$$U(r)\eta_X : X \rightarrow UFX_j$$

makes FX_j free over X . This contradicts to the fact that each endomorphism $f : FX_j \rightarrow FX_j$ with $U(f)\eta_{X_j} = \eta_{X_j}$ is equal to id_{FX_j} . Indeed, since

$$U(r)\eta_X u_j = U(r)UF(u_j)\eta_{X_j} = \eta_{X_j},$$

the image of X in $U(r)\eta_X$ extends X_j to free generators of UFX_j . Then, by interchanging two free generators not belonging to X_j , we get an f above with $f \neq \text{id}_{FX_j}$. \square

4. ABSTRACT ELEMENTARY CLASSES

A class \mathcal{K} of Σ -structures will be always considered as the full subcategory of $\mathbf{Str}(\Sigma)$. The following definition introduces abstract elementary classes using the language of category theory. $|A|$ denotes the cardinality of the underlying set of the Σ -structure A .

Definition 4.1. Let Σ be a finitary signature, \mathcal{K} a class of Σ -structures and \mathcal{M} a class of homomorphisms. Then $(\mathcal{K}, \mathcal{M})$ is an *abstract elementary class* if the following conditions are satisfied:

- (A1) each homomorphism $h : A \rightarrow B$ in \mathcal{M} is an embedding of $A \in \mathcal{K}$ to $B \in \mathcal{K}$ as a submodel,
- (A2) \mathcal{K} -objects together with \mathcal{M} -homomorphisms form a category $\mathcal{K}_{\mathcal{M}}$,
- (A3) $\mathcal{K}_{\mathcal{M}}$ is closed in $\mathbf{Str}(\Sigma)$ under directed colimits,
- (A4) given homomorphisms $g : A \rightarrow B$ and $h : B \rightarrow C$ with $hg, h \in \mathcal{M}$ then $g \in \mathcal{M}$, and
- (A5) there is a cardinal λ such that if $f : A \rightarrow B$ is a submodel embedding with $B \in \mathcal{K}$ then there is $h : A' \rightarrow B$ in \mathcal{M} such that f factorizes through h and $|A'| \leq |A| + \lambda$.

The standard formulation (see [2]) uses colimits of continuous chains in (A3). But it is well known that it implies that it is true for all

directed colimits (see [2], or [1] 1.7; continuous chains are called smooth there). The standard formulation also includes that both \mathcal{K} and \mathcal{M} are closed under isomorphisms. This means that, given K in \mathcal{K} and an isomorphism $h : K \rightarrow X$ with S in $\mathbf{Str}(\Sigma)$ then X belongs to \mathcal{K} and h belongs to \mathcal{M} ; the first property means that \mathcal{K} is *replete* in $\mathbf{Str}(\Sigma)$. But this closedness under isomorphisms is a consequence of (A3). In (A4), we do not need to assume that g is a submodel embedding because it follows from hg being a submodel embedding. Let us recall that submodel embeddings are precisely strong homomorphisms in $\mathbf{Str}(\Sigma)$. This is the standard category theoretic terminology and our reason for avoiding to call \mathcal{M} -morphisms strong embeddings as one does in the abstract elementary classes literature.

We allow many-sorted signatures in 4.1. Since each many-sorted signature can be made single-sorted, this does not change the concept of an abstract elementary class. The following result was also observed in [8] Theorem 5.9 (see [9] Theorem 4.1, as well). There is a related work [6].

Proposition 4.2. *Let $(\mathcal{K}, \mathcal{M})$ be an abstract elementary class. Then $\mathcal{K}_{\mathcal{M}}$ is an LS-accessible category with directed colimits.*

Proof. We observed in 3.5 that $\mathbf{Emb}(\Sigma)$ is an LS-accessible category with directed colimits. There is a regular cardinal κ greater than λ from (A5) such that $\mathbf{Emb}(\Sigma)$ is κ -accessible and κ -presentable objects in $\mathbf{Emb}(\Sigma)$ are precisely Σ -structures A such that $|A| < \kappa$. (A5) then implies that each object of $\mathcal{K}_{\mathcal{M}}$ is a κ -directed colimit of κ -presentable objects in $\mathcal{K}_{\mathcal{M}}$. Thus $\mathcal{K}_{\mathcal{M}}$ is κ -accessible. Moreover, $\mathcal{K}_{\mathcal{M}}$ has an exactly μ -presentable object for each regular cardinal $\kappa \leq \mu$. Thus $\mathcal{K}_{\mathcal{M}}$ is LS-accessible. Following (A5), the embedding of $\mathcal{K}_{\mathcal{M}}$ into $\mathbf{Emb}(\Sigma)$ preserves directed colimits and preserves exactly μ -presentable objects for each regular cardinal $\kappa \leq \mu$. \square

Remark 4.3. (1) Following the proof above, we have $\text{rank}(A) = |A|^+$ for each A in \mathcal{K} with $|A| \geq \lambda$.

(2) $\mathcal{K}_{\mathcal{M}}$ is well accessible (see 3.1).

The following result is closely related to Claim 4.9 in [9]. We just avoid talking about structures and signatures.

Theorem 4.4. *A category \mathcal{L} is equivalent to $\mathcal{K}_{\mathcal{M}}$ for an abstract elementary class $(\mathcal{K}, \mathcal{M})$ if and only if it is an accessible category with directed colimits, whose morphisms are monomorphisms and which admits a nearly full embedding into a finitely accessible category preserving directed colimits and monomorphisms.*

Proof. Necessity is evident (using 4.2). Consider a category \mathcal{L} satisfying the conditions above for a finitely accessible category \mathcal{H} . Let \mathcal{A} be a representative set of finitely presentable objects in \mathcal{H} . Consider the canonical functor

$$E : \mathcal{H} \rightarrow \mathbf{Set}^{\mathcal{A}^{\text{op}}}.$$

(see [1], 1.25). Since objects of $\mathbf{Set}^{\mathcal{A}^{\text{op}}}$ can be viewed as many-sorted unary algebras, $\mathbf{Set}^{\mathcal{A}^{\text{op}}}$ is a full subcategory of $\mathbf{Str}(\Sigma)$ for a finitary signature Σ . Since E is a full embedding, we get a nearly full embedding

$$\mathcal{L} \rightarrow \mathbf{Str}(\Sigma).$$

Since subalgebras are submodels, condition (A1) is satisfied. The same is evidently true for (A2)-(A4). Following 3.2 and 3.1, G is well λ -accessible for some regular cardinal λ . Moreover, we can assume that an object X in $\mathbf{Emb}(\Sigma)$ is μ -presentable if and only if $|X| < \mu$ for each $\mu \geq \lambda$. Let $f : A \rightarrow G(B)$ be a submodel embedding. Then $G(B)$ is μ -presentable in $\mathbf{Emb}(\Sigma)$ for some regular cardinal $\mu \geq \lambda$. Since B is a μ -directed colimit of μ -presentable objects, there is $h : A' \rightarrow B$ in \mathcal{K} such that A' is μ -presentable in \mathcal{K} and f factorizes through h . Since $G(A')$ is μ -presentable in $\mathbf{Emb}(\Sigma)$, $|A'| < \mu < |A| + \lambda$. Thus (A5) is satisfied. \square

Example 4.5. (1) Let T be a theory in a language $L_{\kappa\omega}(\Sigma)$. Then $\mathcal{K} = \mathbf{Mod}(T)$ is an abstract elementary class where \mathcal{M} consists of $L_{\kappa\omega}$ -elementary embeddings. Thus $\mathcal{K}_{\mathcal{M}} = \mathbf{Elem}(T)$.

(2) Let T be a basic theory in a language $L_{\kappa\omega}(\Sigma)$. Then $\mathcal{K} = \mathbf{Mod}(T)$ is an abstract elementary class where \mathcal{M} consists of submodel embeddings. Thus $\mathcal{K}_{\mathcal{M}} = \mathbf{Emb}(T)$.

(3) Let Σ be a single-sorted signature containing just a unary relation symbol P . A simple example of an abstract elementary class \mathcal{K} in Σ which is not closed under $L_{\infty\omega}$ -elementary equivalence (thus not axiomatizable in any $L_{\kappa\omega}(\Sigma)$) is given in [7], 2.10. One takes Σ -structures K such that P_K is countable and the complement of P_K is infinite; \mathcal{M} consists of submodel embeddings which are identities on P . But, $\mathcal{K}_{\mathcal{M}}$ is isomorphic to the category of infinite sets and monomorphisms. The later category is $L_{\omega\omega}$ axiomatizable (in the empty signature). Thus, although that $(\mathcal{K}, \mathcal{M})$ is not axiomatizable in the signature Σ , $\mathcal{K}_{\mathcal{M}}$ is equivalent (as a category) to an axiomatizable one.

Remark 4.6. (1) Shelah's Presentation Theorem (see [2], 4.15) implies that, for each abstract elementary class $(\mathcal{K}, \mathcal{M})$ in a signature Σ , there is a signature $\Sigma \subseteq \Sigma'$ and a basic theory T of $L_{\kappa\omega}(\Sigma')$ such that \mathcal{K} is the full image of the reduct

$$R : \mathbf{Emb}(T) \rightarrow \mathbf{Str}(\Sigma).$$

The reason is that omitting a type of an element means to negate the conjunction of all atomic formulas from this type, which is a basic formula. Moreover, \mathcal{M} is the full image in R of Σ' -submodel embeddings. Since R is finitely accessible, each abstract elementary class is a finitely accessible full image of that from 4.5 (2).

(2) One cannot expect that abstract elementary classes are closed under finitely accessible full images. Let Σ contain a (single-sorted) unary relation symbol U and \mathcal{K} consist of all Σ -structures A such that $|A| = |U_A|$. Since \mathcal{K} is not closed under directed colimits of submodel embeddings, $(\mathcal{K}, \mathcal{M})$ is not an abstract elementary class for \mathcal{M} consisting of submodel embeddings. But $\mathcal{K}_{\mathcal{M}}$ form the full image of

$$R : \mathbf{Emb}(T) \rightarrow \mathbf{Str}(\Sigma)$$

where Σ' also contains a binary operation symbol f and T consists of formulas

$$(\forall x)U(f(x))$$

and

$$(\forall x, y)(f(x) = f(y) \rightarrow x = y).$$

Since R is finitely accessible, $\mathbf{Emb}(T)$ is an abstract elementary class with a finitely accessible full image which is not an abstract elementary class.

5. CATEGORICITY

Definition 5.1. Let λ be an infinite cardinal. A category \mathcal{K} is called *λ -categorical* if it has, up to isomorphism, precisely one object of the presentability rank λ^+ .

Remark 5.2. Following 4.3 (1), the definition 5.1 (suggested in [13]) is in accordance with its model theoretic meaning for abstract elementary classes.

Shelah's Categoricity Conjecture claims that for every abstract elementary class $(\mathcal{K}, \mathcal{M})$ there is a cardinal κ such that if $\mathcal{K}_{\mathcal{M}}$ is κ -categorical then it is λ -categorical for each $\kappa < \lambda$. Following the remark above, this is a statement about the category $\mathcal{K}_{\mathcal{M}}$ – it does not depend on its signature presentation.

Example 5.3. Consider a one-sorted signature Σ given by a sort s and an ω -ary function symbol $f : s^\omega \rightarrow s$. Let T be the $L_{\omega_1\omega_1}(\Sigma)$ theory saying that f is a bijection. Thus T consists of the formula

$$(\forall y)(\exists! x)(f(x) = y)$$

where x is an ω -string of variables, and $\exists!$ denotes unique existence. By general facts about categories of models of $\exists!$ -sentences (called *limit theories*), the category $\mathbf{Mod}(T)$ is locally ω_1 -presentable; see [1], 5.30. Hence, in particular, it is accessible and has directed colimits.

If μ has cofinality ω then

$$\mu^\omega > \mu$$

cf. [5], Corollary 4 of Theorem 17. Thus T does not have models of cardinalities of cofinality ω . Since $\mu^\omega = \mu$ whenever the cardinal μ is of the form ν^ω , $\mathbf{Mod}(T)$ is a large category. Thus there is a proper class of cardinalities in which T has a model and a proper class of cardinalities in which T does not have a model. But, following 3.6, $\mathbf{Mod}(T)$ is LS-accessible. The point is that presentation ranks and cardinalities will never start to coincide.

Example 5.4. Let \mathcal{K} be an accessible category which is not LS-accessible. Let $\mathcal{L} = \mathcal{K} \sqcup \mathbf{Set}$, the disjoint union of \mathcal{K} and the category of sets. \mathcal{L} is an accessible category and there is a proper class of cardinals λ such that \mathcal{L} is λ -categorical and, at the same time, a proper class of cardinals λ such that \mathcal{L} is not λ -categorical.

But we do not know any example of an accessible category with directed colimits which is not LS-accessible. In the same time, we do not know any example of an accessible category with directed colimits, whose morphisms are monomorphisms, and which is not equivalent to $\mathcal{K}_{\mathcal{M}}$ for any abstract elementary class $(\mathcal{K}, \mathcal{M})$. We expect, at least in the second case, that such an example exists. Difficulty in producing it is well illustrated by the next example.

Example 5.5. Let \mathcal{K} be the category of commutative unital C^* -algebras. Gelfand duality says that \mathcal{K} is isomorphic to the dual of the category of compact Hausdorff topological spaces. \mathcal{K} is a locally ω_1 -presentable category. Thus it is an accessible category with directed colimits. There is not known whether \mathcal{K} is equivalent to $\mathbf{Mod}(T)$ where T is a theory of $L_{\kappa\omega}$. P. Bankstone [4] asked whether \mathcal{K} is equivalent to $\mathbf{Mod}(T)$ where T is an $L_{\omega\omega}$ theory of a signature Σ such that $\mathbf{Mod}(T)$ is closed in $\mathbf{Str}(\Sigma)$ under products. A negative answer was independently given in [3] and [14].

REFERENCES

- [1] J. Adámek and J. Rosický, *Locally Presentable and Accessible Categories*, Cambridge University Press 1994.
- [2] J. T. Baldwin, *Categoricity*, Amer. Math. Soc. 2009.

- [3] B. Banaschewski, *More on compact Hausdorff spaces and finitary duality*, Canad. J. Math. 36 (1984), 1113-1118.
- [4] P. Bankston, *Some obstacles to duality in topological algebras*, Canad. J. Math. 34 (1982), 80-90.
- [5] T. Jech, *Set Theory*, Academic Press 1978.
- [6] J. Kirby, *Abstract elementary categories*, preprint 2008.
- [7] D. W. Kueker, *Abstract elementary classes and infinitary logic*, Ann. Pure Appl. Logic 156 (2008), 274-286.
- [8] M. J. Lieberman, *Topological and Category-Theoretic Aspects of Abstract Elementary Classes*, Thesis, The University of Michigan 2009.
- [9] M. J. Lieberman, *Category theoretic aspects of abstract elementary classes*, preprint, April 2010.
- [10] M. Makkai and R. Paré, *Accessible categories: The Foundation of Categorical Model Theory*, Cont. Math. 104, AMS 1989.
- [11] J. Rosický, *Categories of models of languages $L_{\kappa,\lambda}$* , In: Abstr. 9th Winter School on Abstract Analysis, Mathem. Inst. Prague 1981, 153-157.
- [12] J. Rosický, *More on directed colimits of models*, Appl. Cat. Struct. 2 (1994), 71-76.
- [13] J. Rosický, *Accessible categories, saturation and categoricity*, J. Symb. Logic 62 (1997), 891-901.
- [14] J. Rosický, *Elementary categories*, Arch. Math. 52 (1989), 284-288.

T. BEKE

DEPARTMENT OF MATHEMATICS

UNIVERSITY OF MASSACHUSETTS LOWELL

ONE UNIVERSITY AVENUE, LOWELL, MA 01854, U.S.A.

TIBOR_BEKE@UML.EDU

J. ROSICKÝ

DEPARTMENT OF MATHEMATICS AND STATISTICS

MASARYK UNIVERSITY, FACULTY OF SCIENCES

KOTLÁŘSKÁ 2, 611 37 BRNO, CZECH REPUBLIC

ROSICKY@MATH.MUNI.CZ

ABSTRACT ELEMENTARY CLASSES AND ACCESSIBLE CATEGORIES

T. BEKE* AND J. ROSICKÝ**

ABSTRACT. We investigate properties of accessible categories with directed colimits and their relationship with categories arising from Shelah's Abstract Elementary Classes. We also investigate ranks of objects in accessible categories, and the effect of accessible functors on ranks.

1. INTRODUCTION

M. Makkai and R. Paré [10] introduced accessible categories as categories sharing two typical properties of categories of structures described by infinitary first-order theories – the existence of sufficiently many directed colimits and the existence of a set of objects generating all objects by means of distinguished colimits. Their, purely category-theoretical, definition has since then found applications in various branches of mathematics. Very often, accessible categories have all directed colimits. These arise as categories of models and elementary embeddings of infinitary theories with finitary quantifiers. In model theory, S. Shelah went in a similar direction and introduced abstract elementary classes as a formalization of properties of models of generalized logics with finitary quantifiers. Our aim is to relate these two approaches. In Section 5, we introduce a hierarchy of accessible categories with directed colimits. The main result of that section, Corollary 5.7, sandwiches Shelah's Abstract Elementary Classes between two natural families of accessible categories.

Unlike abstract elementary classes, accessible categories are not equipped with canonical 'underlying sets'. Nonetheless, there exists a good substitute for 'size of the model', namely, the *presentability rank* of an object, that can be expressed purely in the language of category theory, i.e. in terms of objects and morphisms. In this sense, accessible categories take an extreme 'signature-free' and 'elements-free' view of

Date: November 12, 2011.

Key words and phrases. abstract elementary class, accessible category.

* Supported by the NSA under Grant H98230-11-1-0172, ** Supported by MSM 0021622409.

abstract elementary classes. From this point of view, Shelah's Categoricity Conjecture, the driving force of abstract elementary classes (see [3]), turns on the subtle interaction between ranks of objects and directed colimits in accessible categories.

The connection between abstract elementary classes and accessible categories was discovered, independently, by M. Lieberman as well; see [8] and [9]. Our Corollary 5.7 simplifies Lieberman's description, [9], Proposition 4.8 and Claim 4.9.

2. ACCESSIBLE CATEGORIES

In order to define accessible categories one just needs the concept of a λ -directed colimit where λ is a regular cardinal number. This is a colimit over a diagram $D : \mathcal{D} \rightarrow \mathcal{K}$ where \mathcal{D} is a λ -directed poset, considered as a category. An object K of a category \mathcal{K} is called λ -presentable if its hom-functor $\text{hom}(K, -) : \mathcal{K} \rightarrow \mathbf{Set}$ preserves λ -directed colimits; here \mathbf{Set} is the category of sets.

A category \mathcal{K} is called λ -accessible, where λ is a regular cardinal, provided that

- (1) \mathcal{K} has λ -directed colimits,
- (2) \mathcal{K} has a set \mathcal{A} of λ -presentable objects such that every object of \mathcal{K} is a λ -directed colimit of objects from \mathcal{A} .

A category is *accessible* if it is λ -accessible for some regular cardinal λ .

A *signature* Σ is a set of (infinitary) operation and relation symbols. These symbols are S -sorted where S is a set of sorts. It is advantageous to work with many-sorted signatures but it is easy to reduce them to single-sorted ones. One just replaces sorts by unary relation symbols and adds axioms saying that there are disjoint and cover the underlying set of a model. Thus the underlying set of an S -sorted structure A is the disjoint union of underlying sets A_s over all sorts $s \in S$. $|A|$ will denote the cardinality of the underlying set of the Σ -structure A . The category of all Σ -structures and homomorphisms (i.e., mappings preserving all operations and relations) is denoted by $\mathbf{Str}(\Sigma)$. A homomorphism is called a substructure embedding if it is injective and reflects all relations. Any inclusion of a substructure is a substructure embedding. Conversely, if $h : A \rightarrow B$ is a substructure embedding then A is isomorphic to the substructure $h(A)$ of B . The category of all Σ -structures and substructure embeddings is denoted by $\mathbf{Emb}(\Sigma)$. Both $\mathbf{Str}(\Sigma)$ and $\mathbf{Emb}(\Sigma)$ are accessible categories, cf. [1], 5.30 and 1.70.

A signature Σ is *finitary* if all relation and function symbols are finitary. For a finitary signature, the category $\mathbf{Str}(\Sigma)$ is locally finitely

presentable and $\mathbf{Emb}(\Sigma)$ is finitely accessible. In both cases, there is a cardinal κ such that, for each regular cardinal $\kappa \leq \mu$, a Σ -structure K is μ -presentable if and only if $|K| < \mu$. This follows from the downward Löwenheim-Skolem theorem; see [10], 3.3.1.

Let λ be a cardinal and Σ be a λ -ary signature, i.e., all relation and function symbols have arity smaller than λ . Given a cardinal κ , the language $L_{\kappa\lambda}(\Sigma)$ allows less than κ -ary conjunctions and disjunctions and less than λ -ary quantifications. A substructure embedding of Σ -structures is called *$L_{\kappa\lambda}$ -elementary* if it preserves all $L_{\kappa\lambda}$ -formulas. A *theory* T is a set of sentences of $L_{\kappa\lambda}(\Sigma)$. $\mathbf{Mod}(T)$ denotes the category of T -models and homomorphisms, $\mathbf{Emb}(T)$ the category of T -models and substructure embeddings while $\mathbf{Elem}(T)$ will denote the category of T -models and $L_{\kappa\lambda}$ -elementary embeddings. The category $\mathbf{Elem}(T)$ is accessible (see [1], 5.42). For certain theories T , the category $\mathbf{Mod}(T)$ does not have μ -directed colimits for any regular cardinal μ and thus fails to be accessible.

A theory T is called *basic* if it consists of sentences

$$(\forall x)(\varphi(x) \Rightarrow \psi(x))$$

where φ and ψ are positive-existential formulas and x is a string of variables. For a basic theory T , the category $\mathbf{Mod}(T)$ is accessible. Conversely, every accessible category is equivalent to the category of models and homomorphisms of a basic theory. All these facts can be found in [10] or [1].

Locally presentable categories are defined as cocomplete accessible categories. Following [1], 1.20, each locally λ -presentable category is locally μ -presentable for each regular cardinal $\mu > \lambda$. Let λ be an uncountable regular cardinal. The category \mathbf{Pos}_λ of λ -directed posets and substructure embeddings is λ -accessible but it is not μ -accessible for all regular cardinals $\mu > \lambda$. Following [10] 2.3, let us write $\lambda \triangleleft \mu$ whenever \mathbf{Pos}_λ is μ -accessible. There are arbitrarily large regular cardinals μ such that $\lambda \triangleleft \mu$ and, at the same time, arbitrarily large regular cardinals μ such that $\lambda \triangleleft \mu$ does not hold. Thus the *accessibility spectrum* of \mathbf{Pos}_λ has a proper class of gaps. Generally, if $\lambda \triangleleft \mu$ then any λ -accessible category \mathcal{K} is μ -accessible; see [10], 2.3.10 or [1], 2.11. By [1], 2.13 (1), one has $\omega \triangleleft \lambda$ for every uncountable regular cardinal λ . Thus a finitely accessible category is μ -accessible for all uncountable regular cardinals μ .

Definition 2.1. We say that a category \mathcal{K} is *well λ -accessible* if it is μ -accessible for each regular cardinal $\lambda \leq \mu$.

\mathcal{K} is *well accessible* if it is well λ -accessible for some regular cardinal λ .

We have just seen that any locally presentable category and any finitely accessible category is well-accessible.

Definition 2.2. Let λ be a regular cardinal. We say that an object K of a category \mathcal{K} has *presentability rank* (or, for brevity, *rank*) λ if it is λ -presentable but not μ -presentable for any regular cardinal $\mu < \lambda$. We will write $\text{rank}(K) = \lambda$.

This concept was introduced by Makkai and Paré under the name of *presentability* of an object. See the very last line of p. 29 of [10].

Remark 2.3. (1) We have mentioned that in $\mathbf{Emb}(\Sigma)$, Σ finitary, there is a cardinal κ such that, for each cardinal $\kappa \leq \mu$, a Σ -structure A is μ -presentable if and only if $|A| < \mu$. Therefore for all large enough structures, $\text{rank}(A) = |A|^+$, i.e., presentability ranks play the role of cardinalities.

(2) Let T be a basic theory in a language $L_{\kappa\omega}(\Sigma)$. Since $\mathbf{Emb}(T)$ is closed under directed colimits in $\mathbf{Emb}(\Sigma)$, $\text{rank}(X) \leq |X|^+$ for each T -model X . By the downward Löwenheim-Skolem theorem, cf. [10] 3.3.1, there is a cardinal μ such that, for $\mu \leq |X|$, we have $\text{rank}(X) = |X|^+$. Moreover, either the presentability ranks of T -models in $\mathbf{Emb}(T)$ form a set or T has models of presentability rank ν^+ for all $\mu \leq \nu$. This means that the *presentability spectrum* of $\mathbf{Emb}(T)$ does not have arbitrarily large gaps.

(3) If T is a basic theory in a language $L_{\kappa\omega}(\Sigma)$ then $\mathbf{Mod}(T)$ does not need to be closed under directed colimits in $\mathbf{Str}(\Sigma)$. But this holds for basic theories T in a language $L_{\kappa\omega}^*(\Sigma)$ allowing only finitary conjunctions. Thus $\mathbf{Mod}(T)$ is an accessible category with directed colimits in this case. [10] calls categories equivalent to such $\mathbf{Mod}(T)$ ∞, ω -*elementary*. Every finitely accessible category is ∞, ω -*elementary*; see [10] 4.3.2. The downward Löwenheim-Skolem theorem applies to ∞, ω -elementary $\mathbf{Mod}(T)$, and implies, similarly to (2), that $\text{rank}(X)$ eventually coincides with $|X|^+$ and the presentability spectrum of $\mathbf{Mod}(T)$ does not have arbitrarily large gaps.

Even if T is only a basic theory in a language $L_{\kappa\omega}(\Sigma)$, then $\mathbf{Mod}(T)$ is closed under λ -directed colimits in $\mathbf{Str}(\Sigma)$ for some regular cardinal λ . Then the downward Löwenheim-Skolem theorem still implies that $\text{rank}(X)$ eventually coincides with $|X|^+$ and the presentability spectrum of $\mathbf{Mod}(T)$ does not have arbitrarily large gaps.

(4) Let T be an arbitrary theory in a language $L_{\kappa\omega}(\Sigma)$. We know that $\mathbf{Elem}(T)$ is an accessible category and, following [1] 5.39, $\mathbf{Elem}(T)$ is closed under directed colimits in $\mathbf{Str}(\Sigma)$. Thus $\mathbf{Elem}(T)$ is an accessible category with directed colimits. Moreover, $\text{rank}(X) = |X|^+$ in

$\mathbf{Elem}(T)$ for all sufficiently large T -models X and the presentability spectrum of $\mathbf{Elem}(T)$ does not have arbitrarily large gaps.

Following [10] 3.2.8, there is a basic theory T' in another language $L_{\kappa\omega}^*(\Sigma')$ such that the categories $\mathbf{Elem}(T)$ and $\mathbf{Mod}(T')$ are equivalent. Thus $\mathbf{Elem}(T)$ is ∞, ω -elementary.

(5) Assuming GCH, in any accessible category \mathcal{K} , $\text{rank}(K)$ is a successor cardinal for all objects K (with the possible exception of a set of isomorphism types). Indeed, let \mathcal{K} be a κ -accessible category and $\text{rank}(K) = \lambda$ with $\kappa < \lambda$. λ is a regular cardinal by definition; if λ was a limit cardinal then (as it is uncountable) it would be a weakly inaccessible cardinal. Since GCH is assumed, λ is inaccessible. Thus, given $\alpha < \kappa < \lambda$ and $\beta < \lambda$, we have $\beta^\alpha < \lambda$. Following [1] 2.13 (4), $\kappa \triangleleft \lambda$. Following [10], 2.3.11, the object K can be exhibited as a κ -directed colimit of κ -presentable objects along a diagram of size less than λ . Let the size of that diagram be ν ; then K is ν^+ -presentable. That would mean $\nu^+ = \lambda$, contradicting that λ is a limit cardinal.

By ‘category’ we always mean a locally small one, i.e., having a set of morphisms between any two objects. A category is called *small* if it has a set of objects. We say that a category is *large* if it is not equivalent to a small category. This means that it has a proper class of non-isomorphic objects. In Remark 2.3 we saw several families of large accessible categories that, starting from some cardinal, possess objects of every possible presentation rank. Since in those examples, this property followed from the downward Löwenheim-Skolem theorem, we will call such categories LS-accessible.

Definition 2.4. An accessible category \mathcal{K} will be called λ -LS-accessible if \mathcal{K} has an object of presentability rank μ^+ for each cardinal $\mu \geq \lambda$.

\mathcal{K} is *LS-accessible* if it is λ -LS-accessible for some cardinal λ .

We will deal with this concept later. However, we have not been able to find any large accessible category which is not LS-accessible.

3. ACCESSIBLE FUNCTORS

A functor $F : \mathcal{K} \rightarrow \mathcal{L}$ is called λ -accessible if \mathcal{K} and \mathcal{L} are λ -accessible categories and F preserves λ -directed colimits. F is called accessible if it is λ -accessible for some regular cardinal λ . By the Uniformization theorem (see [10], 2.5.1 or [1], 2.19), for each accessible functor F there is a regular cardinal λ such that F is λ -accessible and *preserves λ -presentable objects*; it means that if K is λ -presentable then $F(K)$ is λ -presentable. The same is then true for each $\mu \triangleright \lambda$.

Definition 3.1. A functor $F : \mathcal{K} \rightarrow \mathcal{L}$ will be called *well λ -accessible* if \mathcal{K} and \mathcal{L} are well λ -accessible categories and F preserves μ -directed colimits and μ -presentable objects for each $\lambda \leq \mu$. F is called *well accessible* if it is well λ -accessible for some regular cardinal λ .

Remark 3.2. (1) Of course, whenever F preserves λ -directed colimits then it preserves μ -directed ones for $\mu \geq \lambda$.

(2) Every colimit preserving functor between locally λ -presentable categories is well λ -accessible. This immediately follows from the fact that, for regular cardinals $\lambda \leq \mu$, an object of a locally λ -presentable category is μ -presentable if and only if it is a μ -small colimit of λ -presentable objects. Recall that a category is μ -small if it has less than μ morphisms.

(3) Every finitely accessible functor is well finitely accessible. This follows from [1], 2.13 (1) and the fact that, for regular cardinals $\lambda < \mu$, an object of a λ -presentable category is μ -presentable if and only if it is a λ -directed μ -small colimit of λ -presentable objects (see [10], 2.3.11).

(4) Let I be an infinite set and consider the functor $\mathbf{Set} \rightarrow \mathbf{Set}$ that sends X to X^I . This functor is accessible but not well accessible since there are arbitrarily large $|X|$ SUCH that $|X^I| > |X|$, hence the functor does not preserve μ -presentable objects for all arbitrarily large enough μ . On the other hand, for I finite, the functor is well accessible since it is finitely accessible.

A well accessible functor, by definition, will take μ -presentable objects to μ -presentable ones for all large enough μ ; however, it can lower presentability ranks.

Example 3.3. Let $F : \mathbf{Gr} \rightarrow \mathbf{Ab}$ be the abelianization functor, i.e., the reflector from groups to abelian groups. Since F is a left adjoint, it is well finitely accessible by 3.2 (2). There exist simple groups G of arbitrarily large infinite cardinalities. Since $F(G) = 0$ for such a G , 0 is finitely presentable in \mathbf{Ab} and $\text{rank}(G) = |G|^+$, the functor F can lower presentability ranks from κ^+ to ω for arbitrarily large κ .

We say that a functor $F : \mathcal{K} \rightarrow \mathcal{L}$ *reflects λ -presentable objects* if $F(K)$ λ -presentable implies that K is λ -presentable.

Recall that a morphism $g : B \rightarrow A$ is a split epimorphism if there exists $f : A \rightarrow B$ with $gf = \text{id}_A$. Since, in this case, g is a coequalizer of fg and id_B , A is λ -presentable whenever B is λ -presentable (see [1], 1.16).

Definition 3.4. We say that a functor $F : \mathcal{K} \rightarrow \mathcal{L}$ *reflects split epimorphisms* if f is a split epimorphism whenever $F(f)$ is a split epimorphism.

Remark 3.5. Any functor $F : \mathcal{K} \rightarrow \mathcal{L}$ reflecting split epimorphisms is *conservative*, i.e., it reflects isomorphisms. If all morphisms of \mathcal{K} are monomorphisms then F reflects split epimorphisms if and only if it is conservative.

Lemma 3.6. *Let $F : \mathcal{K} \rightarrow \mathcal{L}$ be a λ -accessible functor which reflects split epimorphisms. Then F reflects λ -presentable objects.*

Proof. Let $F(K)$ be λ -presentable in \mathcal{L} . Since \mathcal{K} is λ -accessible, K is a λ -directed colimit $(k_i : K_i \rightarrow K)_{i \in I}$ of λ -presentable objects. Since F preserves λ -directed colimits and $F(K)$ is λ -presentable, there is $i \in I$ and $f : F(K) \rightarrow F(K_i)$ such that $F(k_i)f = \text{id}_{F(K)}$. Since F reflects split epimorphisms, this $k_i : K_i \rightarrow K$ is a split epimorphism. Thus K is λ -presentable. \square

Proposition 3.7. *Let $F : \mathcal{K} \rightarrow \mathcal{L}$ be a well λ -accessible functor which reflects split epimorphisms. Then F preserves presentability ranks μ for $\lambda < \mu$.*

Proof. Let $\text{rank}(K) = \mu$, $\lambda < \mu$. Then $F(K)$ is μ -presentable. Assume that $F(K)$ is ν -presentable for some $\nu < \mu$. Without loss of generality, we can assume that $\lambda \leq \nu$. Following Lemma 3.6, K is ν -presentable, which is a contradiction. Thus $\text{rank}(F(K)) = \mu$. \square

4. ACCESSIBLE CATEGORIES WITH DIRECTED COLIMITS

An important class of accessible categories consists of accessible categories having directed colimits. It includes both ∞, ω -elementary categories and locally presentable ones. On the other hand, the basic $L_{\omega_1 \omega_1}$ theory T of well-ordered sets has both $\mathbf{Mod}(T)$ and $\mathbf{Emb}(T)$ ω_1 -accessible without having directed colimits.

Proposition 4.1. *Any accessible category with directed colimits is well accessible.*

Proof. Let \mathcal{K} be a λ -accessible category with directed colimits and consider a regular cardinal $\lambda < \mu$. Given an object K of \mathcal{K} , there is a λ -directed colimit $(a_i : A_i \rightarrow K)_{i \in I}$ of λ -presentable objects A_i . Let \hat{I} be the poset of all directed subsets of I of cardinalities less than μ (ordered by inclusion). Clearly, \hat{I} is μ -directed. For each $M \in \hat{I}$, let B_M be a colimit of a subdiagram indexed by M . Then B_M is μ -presentable. Since every subset of I having less than μ elements is contained in a directed subset of I having less than μ elements (cf. [1], 2.11), K is a μ -directed colimit of B_M , $M \in \hat{I}$. Thus \mathcal{K} is μ -accessible. \square

Lemma 4.2. *Let \mathcal{K} be a λ -accessible category with directed colimits and K an object of \mathcal{K} which is not λ -presentable. Then $\text{rank}(K)$ is a successor cardinal.*

Proof. Let $\lambda < \mu = \text{rank}(K)$. Following the proof of 4.1, K is a μ -directed colimit of objects K_i which are directed colimits of size $\nu_i < \mu$ of λ -presentable objects. Since K is μ -presentable, it is a retract of some K_i . If $\nu_i < \lambda$ then K_i is λ -presentable and thus K is λ -presentable as well. Let $\lambda \leq \nu_i$. Then K_i is ν_i^+ -presentable and thus K is ν_i^+ -presentable too. Hence $\mu = \nu_i^+$. \square

Proposition 4.3. *Let \mathcal{K} and \mathcal{L} be accessible categories with directed colimits and $F : \mathcal{K} \rightarrow \mathcal{L}$ a functor preserving directed colimits. Then F is well accessible.*

Proof. There is a regular cardinal λ such that both \mathcal{K} and \mathcal{L} are λ -accessible and F preserves λ -presentable objects (see [1] 2.19). Consider a regular cardinal $\lambda < \mu$ and let K be a μ -presentable object of \mathcal{K} . Following the proof of 4.1, K is a μ -directed colimit of objects B_M where each B_M is a directed colimit of less than μ λ -presentable objects. Since K is μ -presentable, it is a retract of some B_M . Since $F(B_M)$ is a directed colimit of less than μ λ -presentable objects, $F(B_M)$ is μ -presentable in \mathcal{L} . Thus $F(K)$ is μ -presentable in \mathcal{L} as a retract of $F(B_M)$. We have proved that F is well λ -accessible. \square

Remark 4.4. Let **Lin** be the category of linearly ordered sets and order preserving injective mappings (they coincide with substructure embeddings). It is a finitely accessible category which is a “minimal” ∞, ω -elementary category in the sense that for every large ∞, ω -elementary category \mathcal{K} there is a faithful functor $E : \mathbf{Lin} \rightarrow \mathcal{K}$ preserving directed colimits; see [10], 3.4.1. Its construction is based on Ehrenfeucht-Mostowski models. Following 4.1 and 4.3, E is well accessible. Since it is faithful, it reflects epimorphisms. Epimorphisms in **Lin** are isomorphisms and thus E reflects split epimorphisms. Following 3.7, E preserves presentability ranks μ starting from some cardinal λ .

Recall that a full subcategory \mathcal{K} of a category \mathcal{L} is called *accessibly embedded* if there is a regular cardinal λ such that \mathcal{K} is closed under λ -directed colimits in \mathcal{L} .

Theorem 4.5. *Accessible categories with directed colimits are precisely reflective and accessibly embedded subcategories of finitely accessible categories.*

Proof. Every reflective and accessibly embedded subcategory of a finitely accessible category \mathcal{L} is accessible (see [1], 2.53) and has directed colimits calculated as reflections of directed colimits in \mathcal{L} . Conversely, let \mathcal{K} be a λ -accessible category with directed colimits. Then \mathcal{K} is equivalent to a full subcategory $\text{Ind}_\lambda(\mathcal{C})$ of $\mathbf{Set}^{\text{cop}}$ consisting of all λ -directed colimits of hom-functors (see [1], 2.26); \mathcal{C} is a small category. Let $\text{Ind}(\mathcal{C})$ be the full subcategory of $\mathbf{Set}^{\text{cop}}$ consisting of all directed colimits of hom-functors. Then $\text{Ind}_\lambda(\mathcal{C})$ is closed in $\text{Ind}(\mathcal{C})$ under λ -directed colimits. Given an object X in $\text{Ind}(\mathcal{C})$, we express it as a directed colimit of hom-functors and take their colimit $F(X)$ in $\text{Ind}_\lambda(\mathcal{C})$ (recall that \mathcal{K} has directed colimits). Clearly, $F(X)$ is a reflection of X in $\text{Ind}_\lambda(\mathcal{C})$. Thus $\text{Ind}_\lambda(\mathcal{C})$ is a reflective subcategory of $\text{Ind}(\mathcal{C})$. \square

Theorem 4.6. *Each large locally presentable category is LS-accessible.*

Proof. Let \mathcal{K} be a locally presentable category. Then \mathcal{K} is locally λ -presentable for some regular cardinal λ and, following [1] 5.30, \mathcal{K} is equivalent to $\mathbf{Mod}(T)$ for a limit theory of $L_{\lambda\lambda}(\Sigma)$ where Σ is an S -sorted signature. Thus it suffices to prove that $\mathbf{Mod}(T)$ is LS-accessible.

Let \mathbf{Set}^S denote the category of S -sorted sets. The category \mathbf{Set}^S is locally finitely presentable and, given an S -sorted set $X = (X_s)_{s \in S}$, $\text{rank}(X) = |X|^+$ where $|X|$ is defined as the cardinality of the disjoint union of X_s , $s \in S$. Let $U : \mathcal{K} \rightarrow \mathbf{Set}^S$ denote the forgetful functor. Since \mathcal{K} is large, there is $t \in S$ such that the sets $U(K)_t$ are arbitrarily large. Let $V : \mathcal{K} \rightarrow \mathbf{Set}$ be the composition of U with the functor $\mathbf{Set}^S \rightarrow \mathbf{Set}$ sending (X_s) to X_t . The functor V preserves limits and λ -directed colimits (see the proof of [1] 5.9). By [1] 1.66, V has a left adjoint F which preserves μ -presentable objects for each $\mu \geq \lambda$, cf. [1] Ex. 1.s(1). Thus $\text{rank}(FX) \leq \text{rank}(X)$ for each X with $\lambda \leq \text{rank}(X)$.

Assume that $\mathbf{Mod}(T)$ is not LS-accessible. Then there is X with $\lambda \leq \text{rank}(X)$ and $\mu = \text{rank}(FX) < \text{rank}(X)$. Since \mathbf{Set}^S is locally μ -presentable, X is a μ -directed colimit of μ -presentable objects X_i , $i \in I$. Let

$$(u_i : X_i \rightarrow X)_{i \in I}$$

denote a colimit cocone. Since F preserves colimits,

$$(Fu_i : FX_i \rightarrow FX)_{i \in I}$$

is a colimit cocone. Since FX is μ -presentable, there is $j \in I$ and $r : FX \rightarrow FX_j$ such that $F(u_j)r = \text{id}_{FX}$. Hence Fu_j is a split epimorphism. There is a mapping $f : X \rightarrow X$ such that $fu_j = u_j$ and $f \neq \text{id}_X$. Since $F(f)UF(u_j) = F(u_j)$, we have $F(f) = \text{id}_{FX}$.

Let $\eta : \text{Id} \rightarrow VF$ be the adjunction unit. Since $\eta_X f = VF(f)\eta_X = \eta_X$, η_X is not a monomorphism. Since the sets VK are arbitrarily large, there is K in \mathcal{K} such that $|X| < |VK|$. Thus there is a monomorphism $X \rightarrow VK$, which implies that η_X is a monomorphism. We get a contradiction. \square

Remark 4.7. (1) We have proved the stronger result that the functor $F : \mathbf{Set} \rightarrow \mathcal{K}$ preserves presentability ranks $\geq \lambda$.

(2) The argument even works for weakly locally presentable categories, i.e., for accessible categories with products. Such categories are equivalent to $\mathbf{Mod}(T)$ where T is a regular theory of $L_{\lambda\lambda}(\Sigma)$ (see [1] Ex. 5.e). The forgetful functor $U : \mathcal{K} \rightarrow \mathbf{Set}^S$ has a weak left adjoint $F : \mathbf{Set}^S \rightarrow \mathcal{K}$ equipped with a natural transformation $\eta : \text{Id} \rightarrow UF$. This follows from the fact that \mathcal{K} is a small-injectivity class in a locally presentable category \mathcal{L} (see [1] 4.8) and thus it is naturally weakly reflective in \mathcal{L} (cf. [2]). This suffices for the argument.

(3) Let \mathcal{K} be a large accessible category with coproducts. Assume that \mathcal{K} is λ -accessible and take the coproduct K of a (representative) set of all λ -presentable objects. Then the functor

$$U = \mathcal{K}(K, -) : \mathcal{K} \rightarrow \mathbf{Set}$$

has arbitrarily large values. U has a left adjoint F given by

$$FX = \coprod_X K$$

By applying the proof of 4.6 to F , we get that \mathcal{K} is LS-accessible.

Example 4.8. Consider a one-sorted signature Σ given by a sort s and an ω -ary function symbol $f : s^\omega \rightarrow s$. Let T be the $L_{\omega_1\omega_1}(\Sigma)$ theory saying that f is a bijection. Thus T consists of the formula

$$(\forall y)(\exists!x)(f(x) = y)$$

where x is an ω -string of variables, and $\exists!$ denotes unique existence. By general facts about categories of models of $\exists!$ -sentences (called *limit theories*), the category $\mathbf{Mod}(T)$ is locally ω_1 -presentable; see [1], 5.30. Hence, in particular, it is accessible and has directed colimits.

If μ has cofinality ω then

$$\mu^\omega > \mu$$

cf. [5], Corollary 4 of Theorem 17. Thus T does not have models of cardinalities of cofinality ω . Since $\mu^\omega = \mu$ whenever the cardinal μ is of the form ν^ω , $\mathbf{Mod}(T)$ is a large category. Thus there is a proper class of cardinalities in which T has a model and a proper class of cardinalities in which T does not have a model. But, following 4.6,

$\mathbf{Mod}(T)$ is LS-accessible. The point is that presentation ranks and cardinalities will never start to coincide.

One can change the theory T to an equational $L_{\omega_1\omega_1}$ theory T' in the signature Σ' consisting of f and ω unary function symbols $g_i : s \rightarrow s$. Equations of T' state that $f : X^\omega \rightarrow X$ and $\langle g_i \rangle_{i \in \omega} : X \rightarrow X^\omega$ are inverse maps (thus, bijections). The functor V of the proof of 4.6 is the underlying set functor $\mathbf{Mod}(T') \rightarrow \mathbf{Set}$ and the functor F is the free algebra functor $\mathbf{Set} \rightarrow \mathbf{Mod}(T')$. Following the proof of 4.6, $\text{rank}(FX) = |X|^+$ for any uncountable set X . For free T -algebras FX , the difference between $\text{rank}(FX)$ and the size of the underlying set $|VFX|$ can become arbitrarily large. This clarifies the last sentence of the preceding paragraph.

Definition 4.9. A functor $H : \mathcal{K} \rightarrow \mathcal{L}$ is *full with respect to isomorphisms* if for any isomorphism $f : HA \rightarrow HB$ there is an isomorphism $\bar{f} : A \rightarrow B$ such that $H(\bar{f}) = f$.

Remark 4.10. Any functor H full with respect to isomorphisms is *essentially injective on objects* in the sense that $HA \cong HB$ implies that $A \cong B$. Any faithful functor full with respect to isomorphisms is conservative, i.e. reflects isomorphisms.

A functor H is called *transportable* if for an isomorphism $f : HA \rightarrow B$ there is a unique isomorphism $\bar{f} : A \rightarrow \bar{B}$ such that $H(\bar{f}) = f$ (this includes $H\bar{B} = B$).

Theorem 4.11. *Let \mathcal{K} be a large accessible category with directed colimits admitting a full with respect to isomorphisms and faithful functor into a finitely accessible category preserving directed colimits. Then \mathcal{K} is LS-accessible.*

Proof. Following 2.3 (2), every finitely accessible category admits a finitely accessible full embedding into $\mathbf{Str}(\Sigma)$ for some finitary signature Σ . Thus we can assume that there is a full with respect to isomorphisms and faithful functor preserving directed colimits $H : \mathcal{K} \rightarrow \mathbf{Str}(\Sigma)$. Consider the pullback

$$\begin{array}{ccc} \mathcal{K} & \xrightarrow{H} & \mathbf{Str}(\Sigma) \\ \bar{G} \uparrow & & \uparrow G \\ \mathcal{L} & \xrightarrow{\bar{H}} & \mathbf{Emb}(\Sigma) \end{array}$$

where G is the inclusion. Since G is transportable, the pullback is equivalent to a pseudopullback (see [10] 5.1.1). Hence \mathcal{L} is accessible

(see [10] 5.1.6) and clearly has directed colimits. The functors \overline{H} and \overline{G} preserve directed colimits, are faithful and full with respect to isomorphisms. In fact, for an isomorphism $f : \overline{G}L_1 \rightarrow \overline{G}L_2$ in \mathcal{K} , there is an isomorphism $\overline{f} : \overline{H}L_1 \rightarrow \overline{H}L_2$ in $\mathbf{Emb}(\Sigma)$ with $G(\overline{f}) = H(f)$. Thus there is an isomorphism $g : L_1 \rightarrow L_2$ with $\overline{G}(g) = f$. The argument for \overline{H} is the same. Consequently, both \overline{H} and \overline{G} are essentially injective on objects (see Remark 4.10) and \overline{G} is surjective on objects. Following 4.1 and 4.3, there is a regular cardinal λ such that both \overline{G} and \overline{H} are well λ -accessible. Since \overline{G} is faithful and full with respect to isomorphisms, it is conservative and thus reflects split epimorphisms (see 4.10 and 3.5).

Now, following 3.7, \overline{G} preserves presentability ranks ν for $\lambda < \nu$. Thus it suffices to prove that \mathcal{L} is λ -LS-accessible. Consider a cardinal $\mu \geq \lambda$. Since \mathcal{K} is large and \overline{G} is essentially injective on objects, \mathcal{L} is large. Since \overline{H} is essentially injective on objects, there is an object L in \mathcal{L} such that $\mu \leq |\overline{H}L|$. Since \mathcal{L} is μ^+ -accessible, L is a μ^+ -directed colimit of μ^+ -presentable objects L_i , $i \in I$. Let $(l_i : L_i \rightarrow L)_{i \in I}$ denote a colimit cocone. There is an embedding $f : A \rightarrow \overline{H}L$ with $|A| = \mu$. Since A is μ^+ -presentable, $f = \overline{H}(l_j)g$ for some $g : A \rightarrow \overline{H}L_j$. Hence $\mu \leq |\overline{H}L_j|$. Since \overline{H} preserves presentability ranks ν for $\lambda < \nu$ (following 3.7), $\overline{H}L_j$ is μ^+ -presentable, i.e., $|\overline{H}L_j| \leq \mu$. Hence $|\overline{H}L_j| = \mu$, i.e., $\text{rank}(\overline{H}L_j) = \mu^+$. Therefore $\text{rank}(L_j) = \mu^+$. We have proved that \mathcal{L} is λ -LS-accessible. \square

Using Ehrenfeucht-Mostowski models, we can prove a stronger result. (Note that any finitely accessible category has a faithful, directed colimit preserving functor into \mathbf{Set} and thus any category \mathcal{K} from 4.11 has this property.)

Theorem 4.12. *Let \mathcal{K} be a large accessible category with directed colimits equipped with a faithful functor $H : \mathcal{K} \rightarrow \mathbf{Set}$ preserving directed colimits. Then \mathcal{K} is LS-accessible.*

Proof. Consider a pullback

$$\begin{array}{ccc} \mathcal{K} & \xrightarrow{H} & \mathbf{Set} \\ \overline{G} \uparrow & & \uparrow G \\ \mathcal{L} & \xrightarrow{\overline{H}} & \mathbf{Emb}(\mathbf{Set}) \end{array}$$

analogous to that in the proof of 4.11. Again, \mathcal{L} is a λ -accessible category with directed colimits. Following 4.5, $\mathcal{L} = \text{Ind}_\lambda \mathcal{C}$ is a full

reflective subcategory of the finitely accessible category $\text{Ind } \mathcal{C}$. Let $F : \text{Ind } \mathcal{C} \rightarrow \mathcal{L}$ be a left adjoint to the inclusion $\mathcal{L} \rightarrow \text{Ind } \mathcal{C}$. The composition $H\overline{G}F : \text{Ind } \mathcal{C} \rightarrow \mathbf{Set}$ preserves directed colimits and thus it is uniquely determined by its domain restriction on \mathcal{C} . Since F is the identity functor on its domain restriction on \mathcal{C} , the domain restriction of $H\overline{G}F$ on \mathcal{C} is faithful. We will prove that $H\overline{G}F$ is faithful. Consider two distinct morphisms $f, g : X \rightarrow Y$ in $\text{Ind } \mathcal{C}$. There are objects $C, D \in \mathcal{C}$ and morphisms $u : C \rightarrow X$, $v : D \rightarrow Y$ and $f', g' : C \rightarrow D$ such that $fu = vf'$ and $gu = vg'$. Hence f' and g' are distinct and thus $H\overline{G}F(f')$ and $H\overline{G}F(g')$ are distinct. Since $H\overline{G}F(v) = G\overline{H}F(v)$ is a monomorphism, $H\overline{G}F(fu)$ and $H\overline{G}F(gu)$ are distinct. Therefore $H\overline{G}F(f)$ and $H\overline{G}F(g)$ are distinct. We have proved that $H\overline{G}F$ is faithful and thus $\overline{G}F$ is faithful.

Since every finitely accessible category is ∞, ω -elementary (see [10] 5.2.6), 4.4 provides a faithful functor $E : \mathbf{Lin} \rightarrow \text{Ind } \mathcal{C}$ preserving directed colimits. Thus $\overline{G}FE$ is faithful and preserves directed colimits. Following 4.4, this functor preserves presentability ranks starting from some cardinal. Thus \mathcal{L} is LS -accessible. In order to prove that \mathcal{K} is LS -accessible, we have to show that \overline{G} is conservative – then one uses 3.5, 4.3 and 3.7. Assume that $\overline{G}(f)$ is an isomorphism. Then $H\overline{G}(f)$, and thus $\overline{H}(f)$ are isomorphisms. Hence f is an isomorphism. \square

Remark 4.13. We have proved that each category \mathcal{K} from 4.12 admits a faithful functor $E : \mathcal{L} \rightarrow \mathcal{K}$ from a finitely accessible category \mathcal{L} which preserves directed colimits and is surjective on objects. Consequently, it admits a faithful functor $\mathbf{Lin} \rightarrow \mathcal{K}$ preserving directed colimits. Moreover, if all morphisms in \mathcal{K} are monomorphisms, we do not need a pullback from the proof of 4.12 and the functor E is then even surjective on morphisms.

We will give an example of an accessible category \mathcal{W} with directed colimits with no faithful functor $\mathcal{W} \rightarrow \mathbf{Set}$ preserving directed colimits. This shows that the two parts of the assumption on the category \mathcal{K} in 4.12 are independent. At the same time, this category \mathcal{W} is an example of an accessible category with directed colimits that is not ∞, ω elementary. On the other hand, we do not know whether the assumptions of 4.12 are necessary for the conclusion. Possibly every accessible category with directed colimits is LS -accessible; possibly every accessible category is LS -accessible.

Example 4.14. Let \mathcal{W} be the category of well-ordered sets where morphisms are either order preserving injective mappings or constant mappings. Since the category \mathcal{W}_0 of well-ordered sets and substructure

embeddings is ω_1 -accessible, \mathcal{W} is ω_1 -accessible as well. The category \mathcal{W}_0 does not have all directed colimits. For example, a countable chain

$$A_0 \rightarrow A_1 \rightarrow \dots A_n \rightarrow \dots$$

such that each A_{n+1} , $0 \leq n$ contains an element a smaller than any $x \in A_n$, does not have a colimit in \mathcal{W}_0 . In fact, it does not have any compatible cocone $A_n \rightarrow A$, $0 \leq n$. In \mathcal{W} , it does have a colimit: the one-element chain 1. One can show that \mathcal{W} has all directed colimits.

Assume that there exists a faithful functor $H : \mathcal{W} \rightarrow \mathbf{Set}$ preserving directed colimits. Following 4.13, there is a faithful functor $\mathbf{Lin} \rightarrow \mathcal{W}$. This is impossible because \mathcal{W} is automorphism rigid – the only isomorphisms $X \rightarrow X$ in \mathcal{W} are identities.

5. ABSTRACT ELEMENTARY CLASSES

Consider the following hierarchy of accessible categories with directed colimits:

- (1) finitely accessible categories,
- (2) ∞, ω -elementary categories,
- (3) accessible categories with directed colimits admitting a faithful functor preserving directed colimits into a finitely accessible category,
- (4) accessible categories with directed colimits.

Each class is contained in the next. The original definition of class (2), ∞, ω -elementary categories, is model-theoretic, as categories equivalent to the category of models and homomorphisms of theories in $L_{\infty, \omega}^*$. There exist categorical descriptions too: these are the categories equivalent to the category of points of some Grothendieck topos; and, by [10] 5.2.6, exactly the 2-categorical limits of finitely accessible categories and finitely accessible functors. Since any finitely accessible category is equipped with a faithful functor into \mathbf{Set} preserving directed colimits, class (3) coincides with the categories from Theorem 4.12. Following 4.14, the inclusion of (3) in (4) is proper. The inclusion of (1) in (2) is also proper. We are not aware of any example of a category belonging to (3) but not to (2). We do know that any category from (2) admits a full embedding preserving directed colimits into a finitely accessible category.

We will show that the categories of models-and-strong-embeddings coming from Shelah’s abstract elementary classes are between (2) and (4). We will also introduce a class of abstract elementary categories as “abstract elementary classes” without any assumption about morphisms being embeddings.

Definition 5.1. A functor $H : \mathcal{K} \rightarrow \mathcal{L}$ will be called *nearly full* if for each commutative triangle

$$\begin{array}{ccc} HA & \xrightarrow{H(h)} & HC \\ & \searrow f & \nearrow H(g) \\ & HB & \end{array}$$

there is $\bar{f} : A \rightarrow B$ in \mathcal{K} such that $H(\bar{f}) = f$.

A subcategory \mathcal{K} of a category \mathcal{L} will be called *nearly full* if the embedding $\mathcal{K} \rightarrow \mathcal{L}$ is nearly full.

Remark 5.2. (1) Every full functor is nearly full. If H is faithful and nearly full, we also have $g\bar{f} = h$ in the definition above. Thus every faithful and nearly full functor reflects split epimorphisms.

(2) [6] calls nearly full subcategories coherent.

(3) Any subcategory \mathcal{K} of a finitely accessible category \mathcal{L} closed under directed colimits is *replete* and full with respect to isomorphisms. In fact, being closed under directed colimits means that each directed colimit in \mathcal{L} of objects from \mathcal{K} belongs to \mathcal{K} . And, directed colimits are determined up to an isomorphism. Recall that replete means to be closed under isomorphic objects.

If \mathcal{K} is a subcategory of an accessible category \mathcal{L} and the embedding of \mathcal{K} to \mathcal{L} preserves directed colimits then the *replete closure* $\bar{\mathcal{K}}$ of \mathcal{K} is closed under directed colimits in \mathcal{L} . Moreover the categories \mathcal{K} and $\bar{\mathcal{K}}$ are equivalent.

Definition 5.3. An accessible categories with directed colimits will be called an *abstract elementary category* if it admits a full with respect to isomorphisms and nearly full embedding preserving directed colimits into a finitely accessible category.

Abstract elementary categories are closely related to Shelah's Abstract Elementary Classes; they differ from those introduced in [6]. We recall the definition of an abstract elementary class using the language of category theory.

Definition 5.4. Let Σ be a finitary signature. A nearly full subcategory \mathcal{K} of $\mathbf{Emb}(\Sigma)$ is called an *abstract elementary class* if it is closed in $\mathbf{Emb}(\Sigma)$ under directed colimits and there is a cardinal λ such that if $f : A \rightarrow B$ is a substructure with $B \in \mathcal{K}$ then there is $h : A' \rightarrow B$ in \mathcal{K} such that f factorizes through h and $|A'| \leq |A| + \lambda$.

The standard formulation (see [3]) uses colimits of continuous chains instead of directed colimits. But it is well known that this does not

change anything (see [3], or [1] 1.7; continuous chains are called smooth there). The standard formulation also includes that \mathcal{K} is replete and full with respect to isomorphisms but 5.2 (3) shows that our formulation is the same. We allow many-sorted signatures in 5.4. Since each many-sorted signature can be made single-sorted, this does not change the concept of an abstract elementary class.

Theorem 5.5. *Let Σ be a finitary signature. A nearly full subcategory \mathcal{K} of $\mathbf{Emb}(\Sigma)$ is an abstract elementary class if and only if it is an accessible category closed under directed colimits in $\mathbf{Emb}(\Sigma)$.*

Proof. Let \mathcal{K} be an abstract elementary class in $\mathbf{Emb}(\Sigma)$. We know that $\mathbf{Emb}(\Sigma)$ is finitely accessible and there is a regular cardinal $\lambda < \kappa$ such that, for each regular cardinal $\kappa \leq \mu$, μ -presentable objects in $\mathbf{Emb}(\Sigma)$ are precisely Σ -structures A such that $|A| < \mu$. Since $\mathbf{Emb}(\Sigma)$ is κ -accessible, Definition 5.4 yields that each object of \mathcal{K} is a κ -directed colimit of κ -presentable objects in \mathcal{K} . Thus \mathcal{K} is κ -accessible.

Conversely, let \mathcal{K} be an accessible nearly full subcategory of $\mathbf{Emb}(\Sigma)$ closed under directed colimits. Then the embedding $\mathcal{K} \rightarrow \mathbf{Emb}(\Sigma)$ is well λ -accessible for some regular cardinal λ greater than κ above. Let $f : A \rightarrow B$ be a substructure embedding with $B \in \mathcal{K}$ and put $\mu = |A| + \lambda$. Since \mathcal{K} is μ^+ -accessible, B is a μ^+ -directed colimit of μ^+ -presentable objects B_i , $i \in I$ in \mathcal{K} . Since the embedding $\mathcal{K} \rightarrow \mathbf{Emb}(\Sigma)$ preserves μ^+ -directed colimits and μ^+ -presentable objects, B is a μ^+ -directed colimit of objects B_i which are μ^+ -presentable in $\mathbf{Emb}(\Sigma)$. Since A is μ^+ -presentable (because $\kappa < \mu^+$), the substructure embedding $f : A \rightarrow B$ factorizes through some B_j , $j \in I$. It suffices to put $A' = B_j$. \square

This result is an improvement of [8] 5.9 and 5.10 (see [9] 4.1 and 4.9, as well; there is a related work [6]). Lieberman assumes that \mathcal{K} is well accessible while we use 4.1 to prove this from the existence of directed colimits. Otherwise, our proof is the same as that of Lieberman.

Remark 5.6. (1) Categories $\mathbf{Elem}(T)$ where T is an $L_{\kappa\omega}$ theory and $\mathbf{Emb}(T)$ where T is a basic $L_{\kappa\omega}$ theory are abstract elementary classes.

(2) Each large abstract elementary class is an LS-accessible category. This follows from 4.11 but also directly from the proof of 5.5. Following this proof, we have $\text{rank}(A) = |A|^+$ for each A in \mathcal{K} with $|A| \geq \lambda$.

We can write our characterization of abstract elementary classes in the language of category theory, i.e., without using Σ -structures.

Corollary 5.7. *A category is equivalent to an abstract elementary class if and only if it is an accessible category with directed colimits whose*

morphisms are monomorphisms and which admits a full with respect to isomorphisms and nearly full embedding into a finitely accessible category preserving directed colimits and monomorphisms.

Proof. Necessity is evident because $\mathbf{Emb}(\Sigma)$ is finitely accessible. Consider a category \mathcal{K} satisfying the conditions above. Let $H : \mathcal{K} \rightarrow \mathcal{L}$ be the corresponding functor into a finitely accessible category. Consider the canonical functor

$$E : \mathcal{L} \rightarrow \mathbf{Set}^{A^{op}}.$$

(see [1], 1.25). E preserves directed colimits (see [1], 1.26). Since objects of $\mathbf{Set}^{A^{op}}$ can be viewed as many-sorted unary algebras, $\mathbf{Set}^{A^{op}}$ is a full subcategory of $\mathbf{Str}(\Sigma)$ for a finitary signature Σ containing operation symbols only. Thus embeddings in $\mathbf{Str}(\Sigma)$ coincide with monomorphisms. Hence the codomain restriction of the composition EH is a full with respect to isomorphisms and nearly full embedding

$$\mathcal{K} \rightarrow \mathbf{Emb}(\Sigma)$$

preserving directed colimits. Following 5.2 (3) and 5.5, \mathcal{K} is equivalent to an abstract elementary class. \square

Remark 5.8. (1) Our definition of abstract elementary category is motivated by Corollary 5.7; it results from dropping the hypotheses on monomorphisms. Any finitely accessible category whose morphisms are not monomorphisms, like posets and isotone mappings, is example of an abstract elementary category that is not an abstract elementary class.

(2) Any abstract elementary class is an abstract elementary category. On the other hand, let \mathcal{K} be an abstract elementary category with a functor H into a finitely accessible category from 5.3. Without any loss of generality, we can assume that $H : \mathcal{K} \rightarrow \mathbf{Str}(\Sigma)$ where Σ is a finitary signature. Consider the pullback from the proof of 4.11. Then \mathcal{L} is an accessible category with directed colimits whose morphisms are monomorphisms and \overline{H} is a full with respect to isomorphisms and nearly full embedding preserving directed colimits and monomorphisms. Thus \mathcal{L} is equivalent to an abstract elementary class. Moreover, the functor \overline{G} is onto on objects and preserves presentability ranks starting from some regular cardinal.

(3) We are not aware of any abstract elementary category which is not ∞, ω -elementary. A simple example of an abstract elementary class \mathcal{K} in Σ which is not closed under $L_{\infty, \omega}$ -elementary equivalence (thus not axiomatizable in any $L_{\kappa\omega}(\Sigma)$) is given in [7], 2.10. There, Σ is a single-sorted signature containing just a unary relation symbol P . Objects of \mathcal{K} are Σ -structures K such that P_K is countable and

the complement of P_K is infinite. Morphisms of \mathcal{K} are substructure embeddings which are identities on P . But, \mathcal{K} is isomorphic to the category of infinite sets and monomorphisms. The latter category is axiomatizable by a basic $L_{\omega\omega}$ theory in the empty signature. Thus it is ∞, ω -elementary.

Proposition 5.9. *Let \mathcal{K} be an abstract elementary category. Then there is an ∞, ω -elementary category \mathcal{L} and a faithful functor $E : \mathcal{L} \rightarrow \mathcal{K}$ which preserves directed colimits and is surjective on objects.*

Proof. Let $\overline{G} : \mathcal{L} \rightarrow \mathcal{K}$ be the embedding of an abstract elementary class \mathcal{L} from 5.8 (2). Shelah's Presentation Theorem (see [3] 4.15) yields a finitary signature $\Sigma \subseteq \Sigma'$, an $L_{\omega\omega}$ -theory T' and a set Γ of quantifier free types in Σ' such that \mathcal{L} consists of Σ -reducts of T' -models omitting types from Γ . Omitting a type can be expressed as an $L_{\infty\omega}$ -sentence. By adding these sentences for types from Γ to T' , we get an $L_{\infty\omega}$ -theory T'' . Since $\mathbf{Elem}(T'')$ is ∞, ω -elementary (see 2.3 (5)), the reduct functor $R : \mathbf{Elem}(T'') \rightarrow \mathcal{L}$ has the desired properties for \mathcal{L} . Thus $\overline{G}R$ has these properties for \mathcal{K} . \square

6. CATEGORICITY

Definition 6.1. Let λ be an infinite cardinal. A category \mathcal{K} is called λ -categorical if it has, up to isomorphism, precisely one object of the presentability rank λ^+ .

Remark 6.2. Following 5.6 (2), the definition 6.1 (suggested in [12]) is in accordance with its model theoretic meaning in abstract elementary classes for sufficiently large cardinals λ .

Shelah's Categoricity Conjecture claims that for every abstract elementary class \mathcal{K} there is a cardinal κ such that \mathcal{K} is either λ -categorical for all cardinals $\kappa \leq \lambda$ or \mathcal{K} is not λ -categorical for any cardinal $\kappa \leq \lambda$. Following 6.2, this is a statement about \mathcal{K} as a category, i.e., it does not depend on the signature in which the abstract elementary class is presented.

By Remark 5.8 (2), Shelah's Categoricity Conjecture is equivalent to the Categoricity Conjecture for abstract elementary categories. It is natural to ask about the status of the Categoricity Conjecture for accessible categories belonging to other levels of our hierarchy. At present, we can offer little information other than this easy observation.

Example 6.3. Suppose \mathcal{K} is an accessible category which is not LS-accessible. Let $\mathcal{L} = \mathcal{K} \sqcup \mathbf{Set}$ be the disjoint union of \mathcal{K} and the category of sets. Then \mathcal{L} is an accessible category and there is a proper class of

cardinals λ such that \mathcal{L} is λ -categorical and, at the same time, a proper class of cardinals λ such that \mathcal{L} is not λ -categorical.

If this \mathcal{K} has directed colimits then the categoricity conjecture fails for the class (4). By 4.12, however, this simple trick does not help for categories in class (3) of our hierarchy.

Remark 6.4. Shelah's conjecture seems to be unknown even for finitely accessible categories, lying at level (1) of the hierarchy. It would be interesting to understand whether the exquisite Galois-theoretic machinery of [3] can be brought to bear implications in this setting.

REFERENCES

- [1] J. Adámek and J. Rosický, *Locally Presentable and Accessible Categories*, Cambridge University Press 1994.
- [2] J. Adámek, H. Herrlich, J. Rosický and W. Tholen, *On a generalized small-object argument for the injectivity subcategory problem*, Cah. Top. Géom. Diff. Cat. XLIII (2002), 83-106.
- [3] J. T. Baldwin, *Categoricity*, Amer. Math. Soc. 2009.
- [4] P. Bankston, *Some obstacles to duality in topological algebras*, Canad. J. Math. 34 (1982), 80-90.
- [5] T. Jech, *Set Theory*, Academic Press 1978.
- [6] J. Kirby, *Abstract elementary categories*, preprint 2008.
- [7] D. W. Kueker, *Abstract elementary classes and infinitary logic*, Ann. Pure Appl. Logic 156 (2008), 274-286.
- [8] M. J. Lieberman, *Topological and Category-Theoretic Aspects of Abstract Elementary Classes*, Thesis, The University of Michigan 2009.
- [9] M. J. Lieberman, *Category theoretic aspects of abstract elementary classes*, Annals Pure Appl. Logic 162 (2011), 903-915.
- [10] M. Makkai and R. Paré, *Accessible categories: The Foundation of Categorical Model Theory*, Cont. Math. 104, AMS 1989.
- [11] J. Rosický, *More on directed colimits of models*, Appl. Cat. Struct. 2 (1994), 71-76.
- [12] J. Rosický, *Accessible categories, saturation and categoricity*, J. Symb. Logic 62 (1997), 891-901.

T. BEKE

DEPARTMENT OF MATHEMATICS

UNIVERSITY OF MASSACHUSETTS LOWELL

ONE UNIVERSITY AVENUE, LOWELL, MA 01854, U.S.A.

TIBOR_BEKE@UML.EDU

J. ROSICKÝ

DEPARTMENT OF MATHEMATICS AND STATISTICS

MASARYK UNIVERSITY, FACULTY OF SCIENCES

KOTLÁŘSKÁ 2, 611 37 BRNO, CZECH REPUBLIC

ROSICKY@MATH.MUNI.CZ